



1655

PATENT #16

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

RECEIVED Jm

MAY 22 2003

TECH CENTER 1600/2900 S/27/0

In re Application of: Pype et al.

Serial No.: 09/697,863

Filed: October 27, 2000

For: CD40-INTERACTING AND TRAF-
INTERACTING PROTEINS

Confirmation No.: 7540

Examiner: S. Liu

Group Art Unit: 1653

Attorney Docket No.: 2676-4555US

CERTIFICATE OF MAILING

I hereby certify that this correspondence along with any attachments referred to or identified as being attached or enclosed is being deposited with the United States Postal Service as First Class Mail on the date of deposit shown below with sufficient postage and in an envelope addressed to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

May 16, 2003
Date

Signature

Betty Vowles
Name (Type/Print)

COMMUNICATION

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

Enclosed is a certified copy of Priority Document 98201392.2 filed April 29, 1998, for the above-referenced case.

Respectfully submitted,

Bretton L. Crockett
Registration No. 44,632
Attorney for Applicant(s)
TRASKBRITT, P.C.
P.O. Box 2550
Salt Lake City, Utah 84110-2550
Telephone: 801-532-1922

Date: May 15, 2003
BLC/bv



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

RECEIVED

MAY 22 2003

TECH CENTER 1600/2900



Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

98201392.2

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

DEN HAAG, DEN
THE HAGUE,
LA HAYE, LE

22/04/02



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

**Blatt 2 der Beschreibung
Sheet 2 of the certificate
Page 2 de l'attestation**

Anmeldung Nr.:
Application no.:
Demande n°: 98201392.2

Anmeldetag:
Date of filing:
Date de dépôt: 29/04/98

Anmelder:
Applicant(s):
Demandeur(s):
Vlaams Interuniversitair Instituut voor Biotechnologie vzw.
9052 Zwijnaarde
BELGIUM

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:
Novel CD40 interacting proteins

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:
Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

/

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing:
Etats contractants désignés lors du dépôt:

AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE

Bemerkungen:
Remarks:
Remarques:

Printed:22-04-2002

+32 9 2

SPEC

98201392

OAV7 / SPY / CRAP / 001- 018 / EP

NOVEL CD40 INTERACTING PROTEINS

The invention relates to novel CD40 binding proteins, which can be used as modulators of the CD40 signalling pathway and/or the CD40-induced Nuclear factor kappa B (NF- κ B) activating pathway and are thus useful in the treatment of CD40 related diseases (e.g. inflammatory diseases) and/or NF- κ B related diseases and/or in the improvement of anti-tumour treatments.

The invention also relates to nucleic acids coding for said novel CD40 interacting proteins.

The invention relates further to the use of polypeptides derived from these CD40 interacting proteins in the treatment of CD40 and/or NF- κ B related diseases and/or cancer.

Furthermore, the invention concerns pharmaceutical preparations comprising the novel CD40 interacting proteins or polypeptides derived from these proteins.

CD40 is a receptor of the TNF- receptor superfamily (Banchereau *et al.*, 1994), which is expressed at the surface of B-cells, antigen presenting cells (APC), and several non-haematopoietic cells such as endothelial cells (Hollenbaugh *et al.*, 1995), epithelial cells (Galy & Spits, 1992), fibroblasts (Fries *et al.*, 1995) and keratinocytes (Gaspari *et al.*, 1996). The ligand for CD40, CD40L, occurs mainly on activated T-cells. Up to now, the role of CD40 was mainly studied in the context of the T-cell APC / B-cell interaction (for a review, see Noelle, 1996). Amongst others, the CD40-CD40L interaction seems to be important for the T-cell mediated immunity and for primary and secondary humoral immune responses. These findings were confirmed by experiments in mouse models, where one was able to show that

treatment with anti-CD40L antibodies resulted in blocking of the development of mouse equivalents of human autoimmune diseases such as arthritis (Durie *et al.* 1993), oophoritis (Griggs *et al.*, 1996) and multiple sclerosis (Gerritse *et al.*, 1996).

- 5 Activation and transduction through the CD40 pathway is in a large part responsible for B cell activation and accordingly, the humoral immune response.

Apart from NF- κ B, factors that can be activated by stimulation of CD40 are NF-AT (Francis *et al.*, 1995) c-Jun, ATF-2 and IRF-1 (Karmann *et al.*, 1996).

- 10 All these factors play an important role in inflammation.

The CD40L induced signal transduction is, as for the case of TNF, mediated by the binding of TNF-Receptor Associated Factors (TRAF's) to the cytoplasmic domain of the receptor. Chaudhuri *et al.* (1997) demonstrated that, at least in human B cell lines, CD40 and TRAF2 are constitutively

- 15 associated with each other, and that this association is inhibited by CD40 mediated signals. Apart from the binding with TRAF 2, the cytoplasmic domain of CD40, which consists of 62 amino acids at positions 196-257 (mature human CD40 - numbering according to Kashiwada *et al.*, 1998), is known to associate with TRAF3, TRAF5, TRAF6 and Janus kinase 3. TRAF 6
- 20 binds to the amino-terminal cytoplasmic tail of CD40 at positions 210-225, although the possibility can not be excluded that full association of TRAF6 with CD40 may also require the carboxy-terminal part at positions 226-249 (Ishida *et al.*, 1996). TRAF 2, TRAF3 and TRAF5 bind to the carboxy-terminal CD40 cytoplasmic domain at positions 226-249 (Ishida *et al.*, 1996).

Stimulation of CD40 results in activation of protein kinases, NF- κ B, the mitogen-activated protein kinase and Janus kinase 3 / signal transducer and activator of Transcription 3. Moreover, stimulation of CD40 mediates critical biological effects in B cell growth, survival and differentiation.

- 5 It is known that TRAF2 and TRAF5 play a role in NF- κ B activation in signalling through CD40, as well as TNF-RI, TNF-RII, CD30 and lymphotoxin β receptor. TRAF6 participates in NF- κ B activation signalled by CD40 and IL-1 receptor.

In addition to these data in WO 96/16665 and WO 96/28588 are disclosed a
10 TRAF like protein that binds to the cytoplasmic domain of CD40.

Surprisingly, it is shown in this invention that two other proteins exist interacting with the cytoplasmic domain of CD40. Even more surprisingly, none of these proteins shows significant homology with one of the known CD40 interacting proteins, neither is there homology between the two proteins
15 themselves. These proteins should therefore be considered as two new classes of CD40 interacting proteins

One aspect of this invention is to offer said novel proteins to modulate and/or inhibit CD40 signalling and/or CD40-induced NF- κ B activation.

One embodiment of the invention is a protein with SEQ ID NO.2. Another
20 embodiment of the invention is a protein with SEQ ID NO.4. A further embodiment of the invention concerns a protein with SEQ ID NO.6.

A further aspect of the current invention is the use of above mentioned proteins, or biologically active fragments of these proteins, to modulate and/or inhibit CD40 signalling and/or CD40-induced NF- κ B activation.

Another aspect of the invention is the use of above mentioned proteins or biologically active fragments of these proteins to screen for compounds that interfere in the interactions of said proteins or fragments with other compounds of the CD40 related signalling pathway.

- 5 Another aspect of the invention consists of DNA molecules encoding for the above mentioned proteins.

The invention also relates to a pharmaceutical composition comprising one or more of the above mentioned proteins or fragments in a biologically active amount for the treatment of CD40 and/or NF- κ B related diseases such as
10 atherosclerosis, arthritis, multiple sclerosis, systemic lupus erythematosus, graft rejection and the like.

In another aspect the present invention relates to a pharmaceutical composition comprising one or more compounds obtainable by the above mentioned screening method for the treatment of CD40 and/or NF- κ B related
15 diseases such as atherosclerosis, arthritis, multiple sclerosis, systemic lupus erythematosus, graft rejection and the like.

BRIEF DESCRIPTION OF THE FIGURES

- 20 **Figure 1:** schematical representation of CRAP (=CD40 receptor associated protein) and the deletion mutants of CRAP used in two hybrid screening assays. The deletion mutants consist of the following amino acids of the original CRAP sequence: 54 to 362 (4F2), 54 to 273 (4F2d3), 54 to 236 (4F2d2) and 54 to 140 (4F2d1).

Figure 2: Northern blot analysis of (a) human tissue, using a human CRAP probe; (b) adult mouse tissue, using a mouse CRAP probe; (c) embryonic mouse tissue, using a mouse CRAP probe. The hybridization of GAPDH is used as a control.

5

DEFINITIONS

The following definitions are provided in order to illustrate and define the meaning and scope of the various terms used in the current description.

The term "treatment" or "treating" or "treat" means any treatment of a disease
10 in a mammal, including : (1) preventing the disease, that is, causing the clinical symptoms of the disease not to develop; (2) inhibiting the disease, that is, arresting the development of the clinical symptoms; and/or (3) relieving the disease, that is, causing the regression of clinical symptoms.

The term "effective amount" means a dosage sufficient to provide treatment
15 for the disease state being treated. This will vary depending on the patient, the disease and the treatment being effected.

"Capable to interact" means that a protein can form a complex with another protein, as can be measured using a yeast two-hybrid system, or with co-immunoprecipitation, or with equivalent systems known to people skilled in
20 the art.

"Functional" protein or fragment means a protein or fragment that is capable to interact with the cytoplasmic part of CD40, or with another protein of the CD40 and/or NF- κ B related pathway.

"Homology to TRAF-proteins" means that the typical structural features found in the current TRAF proteins (TRAF1 - TRAF6) are present. These features comprise a RING finger motif at the amino terminus followed by five or more zinc fingers and a so-called TRAF domain known to a person skilled in the art.

- 5 The "cytoplasmic part of CD40" means a part comprising the 62 carboxy terminal amino acids of human CD40 (amino acid 216-277; Stamenkovic *et al.* 1989), or the homologous mouse sequence, or another homologous sequence with a similar biological activity.

- 10 "Nucleic acid" means genomic DNA, cDNA, double stranded or single stranded DNA, messenger RNA or any form of nucleic acid known to the people skilled in the art.

- "Compound" means any chemical or biological compound, including simple or complex inorganic or organic molecules, peptides, peptido-mimetics, proteins, antibodies, carbohydrates or nucleic acids, that interferes with the interaction
15 of a protein depicted in SEQ ID NO. 2, 4 or 6 with a compound of the CD40 and/or NF- κ B related pathway.

EXAMPLES

- 20 **Example 1: isolation of the CD40 interacting proteins**

Yeast two-hybrid screening.

The two-hybrid screening was performed by the interaction trap cloning method, which is often referred to as the LexA two-hybrid system (Gyuris *et*

al., 1993). The DNA encoding the cytoplasmic part of CD40 (62 amino acids, from residue 216 to 277, where the open reading frame ends, according to the sequence and numbering as given in Stamenkovic *et al.* (1989)) was generated by PCR and inserted into the EcoRI-Sall digested pEG202 vector (Gyuris *et al.*, 1993), in frame with the LexA DNA-binding domain (hereinafter the "bait plasmid"). Screening was performed using a HeLa cell fusion library in the plasmid pJG45 (hereinafter the "prey plasmid"), that was obtained from the laboratory of R. Brent (Harvard Med. School, Boston, MA, USA). Transformation of EGY48 yeast (MAT alpha, *his3*, *trp1*, *ura3-52*, *leu2::pLeu2-LexAop6*) with the prey plasmid, the bait plasmid and the p8op-LacZ (Clontech) reporter plasmid was performed by the Lithium Acetate transformation method (Gietz *et al.*, 1995). The two-hybrid screening was conducted as described in the manual distributed by the laboratory of R. Brent (published in "Gene probes- A practical approach, Oxford University press").

Results of the two-hybrid screening.

Yeast containing bait plasmid and lacZ reporter plasmid was transformed with 20 microgram prey library plasmid and plated on glucose medium lacking tryptophan, histidine and uracil, to select for the presence of all three plasmids. In total, approximately 1.5×10^6 transformants were obtained. The transformants were harvested and frozen at -70°C in a glycerol solution (65% v/v glycerol; 0.1 M MgSO_4 , 25 mM Tris pH 7.4). From these stocks, 20×10^6 colony forming units were plated on galactose medium lacking leucine,

tryptophan, histidine and uracil, to screen for protein-protein interaction. Yeast colonies growing on the latter selective medium were further checked for interaction by screening for blue/white staining on medium containing X-gal and galactose. The colonies displaying the following phenotype were picked
5 for further analysis: i) no growth on glucose containing medium lacking leucine, ii) growth on galactose containing medium lacking leucine, iii) white on medium containing glucose and X-gal, iv) blue on medium containing galactose and X-gal.

Plasmids were isolated from the yeast with the proper phenotype. Analysis of
10 the obtained prey plasmids revealed that the entire screening had finally resulted in the isolation of three different cDNA inserts. Sequencing of the clones showed that, in addition to a partial cDNA for TRAF3, we had isolated two novel cDNA's, termed CRAP and 4C4.

15 Isolation of the full length cDNA

Full length human CRAP cDNA was obtained by screening a HUVEC cDNA library, made in the laboratory, with the CRAP fragment as probe. A cDNA of about 2 kb was isolated, with an open reading frame of 1086 nucleotides long, encoding for a protein of 362 amino acids long (SEQ ID NO.2).

20 The mouse CRAP homologue was obtained by screening the EST database and aligning the homologous sequences. Human and mouse CRAP are approximately 65% identical and 70% similar on the amino acid level. The mouse sequence is shown in SEQ ID NO. 3.

Example 2: sequence analysis of the cDNA's

Nucleotide sequence analysis was carried out using dy terminator mix and a 310 Genetic analyzer from Perkin Elmer. The sequence of CRAP is shown in SEQ ID NO.1 whereas the sequence of 4C4 is shown in SEQ ID NO.5.

- 5 The CRAP sequence shows a low homology (30% similarity at amino acid level) with Nocturin, a protein that is expressed in the photoreceptor of the eye of *Xenopus laevis* (Green and Beshare, 1996). The partial sequence of the mouse homologue of Nocturin is also known (Puech *et al.*, 1997). Additionally, there is some homology with EST sequences (e.g. genbank EST
- 10 c23016, aa162513, aa571061, t87026, h45114, aa196281, h94108 and aa337396) and with the C-terminal part of the yeast transcription factor *CCR4* (Malvar *et al.*, 1992). All these homologies are low, and it is clearly unexpected that a human homologue of these proteins would bind to the cytoplasmic domain of CD40.
- 15 It is interesting to note that, unexpectedly, CRAP neither 4C4 show any significant homology with TRAF's or other proteins known to interact with CD40.

20 **Example 3: study of the interaction of CRAP protein, 4C4 protein and CRAP protein fragments with other proteins using a yeast two-hybrid interaction assay**

The potential binding of CRAP to other proteins was assessed using the yeast two-hybrid assay. The experimental outline is similar to the one described for the two-hybrid screening. However, here the plasmids for bait,

prey and lacZ report r were transformed simultaneously into the EGY48 yeast strain. Positive interaction was determined either by the growth ph notyp (growth on medium lacking leucine in th presence of galactose, and not in the presence of glucose) or by the blue/white staining on X-gal containing plates (blue colonies only on galactose containing plates, not on glucose containing plates). cDNA's for TRAF2 and for the cytoplasmic regions of CD30, CD40 and TNF-RII were generated by PCR using the pfu polymerase (Promega). PCR fragments encoding RIP, TRADD and FADD were cloned in pCDNA3 (Invitrogen, Carlsbad, CA). cDNA of TRAF3 was obtained from the laboratory of Dixit, Dept Pathol., Univ. Michigan Med. School, MI, USA). The color formation was evaluated as strong and fast (++), strong but slow (+), weak and slow (+/-), none (-) or not determined (nd)

The results for CRAP protein and CRAP fragments are summarized in Table I and Figure 1.

15

Table I

	CRAP	4F2	4F2d3	4F2d2	4F2d1	4C4	—
CD40	++	++	+/-	+/-	+/-	+	-
CD30	++	++	+/-	+/-	+/-	+	-
TNF-RII	+	+	-	-	-	+	-
LMP-1	-	-	nd	nd	nd	-	-
TRAF2	-	-	nd	nd	nd	nd	-
TRAF3	+	+	-	-	-	nd	-
RIP	++	++	+/-	+/-	+/-	nd	+/-
TRADD	+	nd	nd	nd	nd	nd	-
FADD	-	nd	nd	nd	nd	nd	-
4F2	++	++	-	-	-	+	-
4C4	++	++	-	-	-	+	-

CRAP, as well as the longest CRAP fragment (aa 54 - 362) shows a strong interaction with CD40, CD30, RIP and with 4C4, and a weaker interaction with TNF-RII and TRAF3. Remarkably, CRAP can also self-associate. CRAP fragments, missing the C-terminal end (aa 274 - 362) show only a weak interaction.

4C4 protein is interacting with CD40, CD30, TNF-RII, with the longest fragment of CRAP and with a deletion mutant of TRAF3 which still contains the largest part of the TRAF domain (from aa 380 to the carboxy terminal end of the protein. A smaller form of 4C4 (from amino acid 2 - amino acid 245 in SEQ ID NO.6) is also capable to interact with CD40.

Example 4: expression pattern of CRAP and 4C4

The CRAP gene is widely expressed, as was already indicated by the presence of several partial CRAP cDNA's in the EST sequence data base. The CRAP expression was analyzed by Northern blot analysis against mRNA from different tissues, both from human and mouse (Figure 2). Human CRAP is present as a 2.2 kb transcript in all tissues tested. Besides the 2.2 kb transcript, there is an additional 1.7 kb transcript in a testis sample. (Figure 2A).

Human CRAP expression was further tested and found in the B-cell lines BJAB (Menezes *et al.*, 1975) and DG75 (Ben-Bassat *et al.*, 1977), in the Jurkat T-cell line and in HUVECs.

For mouse CRAP, two transcripts, one of 2.2 kb and one of 3.8 kb can be found on a murine multiple Northern blot (Figure 2B). Mouse CRAP mRNA is also detected in all tissues tested, but to a lower extent in skeletal muscle. Both mouse transcripts are not only present in adult animals, but can also be detected in mouse embryo's 7 and 17 days post coitum. These results are an indication that CRAP plays an important role in early development.

On a multiple tissue Northern blot, a 4C4 probe recognizes 3 transcripts, of 1.6kb, 3.5 kb and 7.5kb. All three mRNA's are present in spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes. The expression of the 3.5 kb transcript is most prominent in testis. In ovary, the signal of the 7.5 kb mRNA is strongest.

REFERENCES

- 15 Banchereau, J.; Bazan, F.; Blanchard, D.; Briere, F.; Galizzi, J.P.; Van Kooten, C.; Liu, Y.J.; Rousset, F.; Saeland, S. (1994). The CD40 antigen and its ligand. *Annu. Rev. Immunol.* 12, 881
- 20 Ben-Bassat, H.; Goldblum, N.; Mitrani, S.; Goldblum, T.; Yoffey, J.M.; Cohen, M.M.; Bentwich, Z.; Ramot, B.; Klein, E.; Klein, G. (1977) Establishment in continuous culture of a new type of lymphocyte from a "Burkitt like" malignant lymphoma (line D.G.-75). *Int. J. Cancer* 19, 27.
- 25 Chaudhuri, A.; Orme, S.; Eilan, S.; Cherayl, B.J. (1997). CD40 mediated signals inhibit the binding of TNF receptor associated factor 2 to the CD40 cytoplasmic domain. *J. Immunol.* 159, 4244.
- 30 Durie, F.H.; Fava, R.A.; Foy, T.M.; Aruffo, A.; Ledbetter, J.A.; Noelle, R.J. (1993). Prevention of collagen-induced arthritis with an antibody to gp39, the ligand for CD40. *Science* 261, 1328.
- Francis, D.A.; Karras, J.G.; Ke, X.Y.; Sen R.; Rothstein, T.L. (1995). Induction of the transcription factors NF-kappa B, AP-1 and NF-AT during B cell stimulation through the CD40 receptor. *Int. Immunol.* 7, 151.

Fries, K.M.; Sempowski, G.D.; Gaspari, A.A.; Blieden, R.J.; Looney, R.J.; Philips, R.P. (1995). CD40 expression by human fibroblasts. *Clin. Immunol. Immunopath.* 77, 42.

- 5 Galy, A.; Spits, H. (1992). CD40 is functionally expressed on human thymic epithelial cells. *J. Immunol.* 149, 775.

- Gaspari, A.A.; Sempowski, G.D.; Chess, P.; Gish, J.; Phillips, R.P. (1996). Human epidermal keratinocytes are induced to secrete interleukin-6 and costimulate T lymphocyte proliferation by a CD40-dependent mechanism. *Eur. J. Immunol.* 26, 1371.

- 10 Gerritse, K.; Laman, J.D.; Noelle, R.J.; Aruffo, A.; Ledbetter, J.A.; Boersma, W.J.A.; Claassen, E. (1996). CD40-CD40 ligand interactions in experimental allergic encephalomyelitis and multiple sclerosis. *Proc. Natl. Acad. Sci. USA* 93, 2499.

- Gietz, R.D.; Schiestl, R.H.; Willems, A.R.; Woods, R.A. (1995). Studies on the transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure. *Yeast*, 11, 355.

- 20 Green, C.B.; Beshare, J.C. (1996). Identification of a novel vertebrate circadian clock-regulated gene encoding the protein nocturnin. *Proc. Natl. Acad. Sci. USA* 93, 14884.

- 25 Griggs, N.; Agersborg, S.; Noelle, R.J.; Ledbetter, J.; Linsley, P.; Tung, K. (1996). The relative contribution of the CD28 and gp39 costimulatory pathways in the clonal expansion and pathogenic acquisition of self-reactive T cells. *J. Exp. Med.* 180, 801.

- 30 Gyuris, J.; Golemis, E.; Chertkov, H.; Brent, R. (1993). Cdi1, a human G1 and S phase protein phosphatase that associates with Cdk2. *Cell* 75, 791.

- 35 Hollenbaugh, D.; Mischel-Petty, N.; Edwards, C.P.; Simon, J.C.; Denfeld, R.W.; Kienem, P.A.; Aruffo, A. (1995). Expression of functional CD40 by vascular endothelial cells. *J. Exp. Med.* 182, 33.

- 40 Ishida, T.; Tojo, T.; Aoli, T.; Kobayashi, N.; Ohishi, T.; Watanabe, T.; Yamamoto, T.; Inoue, J. (1996). TRAF5, a novel tumor necrosis factor receptor associated factor family protein, mediated CD40 signalling. *Proc. Natl. Acad. Sci. USA* 93, 9437.

- 45 Karmann, K.; Min, W.; Fanslow, W.C.; Prober, J.S. (1996) Activation and homologous desensitization of human endothelial cells by CD40 ligand, tumor necrosis factor and interleukin 1. *J. Exp. Med.* 184, 173.

Kashiwada, M.; Shirakata, Y.; Inoue, J.-I.; Nakano, H.; Okazaki, K.; Okumura, K.; Yamamoto, T.; Nagaoka, H.; Takemori, T. (1998) Tumor Necrosis Factor

Receptor-associated Factor 6 (TRAF6) stimulates extracellular signal-regulated kinase (ERK) activity in CD40 signalling along a Ras-Ind p ndent pathway. *J. Exp. Med.* 187, 237.

- 5 Malvar, T.; Biron, R.W.; Kaback, D.B.; Denis, C.L. (1992) The CCR4 protein from *Saccharomyces cerevisiae* contains a leucine rich repeat region which is required for its control of *ADH2* gene expression. *Genetics*, 132, 951.
- 10 Menezes, J.; Leibold, W.; Klein, G.; Clements, G. (1975). Establishment and characterization of an Epstein-Barr virus (EBV)-negative lymphoblastoid B cell line (BJA-B) from an exceptional, EBV-genome-negative African Burkitt's lymphoma. *Biomedicine* 22, 276.
- 15 Noelle, R.J. (1996). CD40 and its ligand In host defense. *Immunity*, 4, 415-419.
- Puech, A.; Dupressoir, A.; Loireau, M.; Mattei, M.; Heidemann, T. (1997). Characterization of two age-induced intracisternal A-particle-related transcripts In the mouse liver. *J. Biol. Chem.* 272, 5995.
- 20 Stamenkovic, I.; Clark, E.A.; Seed, B. (1989) A B-lymphocyte activation molecule related to the nerve growth factor receptor and induced by cytokines in carcinomas. *Embo J.* 8, 1403.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

10

(A) NAME: Vlaams Interuniversitair Instituut voor
Biotechnologie
(B) STREET: Rijvisschestraat 118 box 1
(C) CITY: Zwijnaarde
(E) COUNTRY: Belgium
(F) POSTAL CODE (ZIP): 9052B
(G) TELEPHONE: +32 9 2446611
(H) TELEFAX: +32 9 2446610

15

(ii) TITLE OF INVENTION: Novel CD40 interacting proteins

(iii) NUMBER OF SEQUENCES: 6

20

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

25

(2) INFORMATION FOR SEQ ID NO: 1:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1920 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

40

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

45

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION:20..1108

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GTGCAGAGGC GGCAGGAAGA TGCAGTTGGG GAGTTGCCTG GAGGGCGGGA GCGAGGCGGC
60

55

GGAGGAAGAG GCGGAGCCTG AGGTGAAAAA GCGGCGACTT CTGTGTGTGG AGTTTGCCTC
120

GGTCGCAAGC TCGGATGCCG CAGTGGCTCA GTGCTTCCTG GCCGAGAACG ACTGGGAGAT
180

GGAAAGGGCT CTGAACTCCT ACTTCGAGCC TCCGGTGGAG GAGAGCGCCT TGGAAACGCCG
240

5 ACCTGAAACC ATCTCTGAGC CCAAGACCTA TGTTGACCTA ACCAATGAAG AAACAACCTGA
300

TTCCACCACT TCTAAAATCA GCCCATCTGA AGATACTCAG CAAGAAAATG GCAGCATGTT
360

10 CTCTCTCATT ACCTGGAATA TTGATGGATT AGATCTAAAC AATCTGTCAG AGAGGGCTCG
420

AGGGGTGTGT TCCTACTTAG CTTTGTACAG CCCAGATGTG ATATTTCTAC AGGAAGTTAT
480

15 TCCCCCATAT TATAGCTACC TAAAGAAGAG ATCAAGTAAT TATGAGATTA TTACAGGTCA
540

20 TGAAGAAGGA TATTTACAG CTATAATGTT GAAGAAATCA AGAGTGAAAT TAAAAGCCA
600

AGAGATTATT CCTTTTCCAA GTACCAAAT GATGAGAAAC CTTTTATGTG TGCATGTGAA
660

25 TGTGTCAGGA AATGAGCTTT GCCTTATGAC ATCCCATTTG GAGAGCACCA GAGGGCATGC
720

TGCGGAACGA ATGAATCAGT TAAAATGGT TTAAAGAAA ATGCAAGAGG CTCCAGAGTC
780

30 AGCTACAGTT ATATTTGCAG GAGATACAAA TCTAAGGGAT CGAGAGGTTA CCAGATGTGG
840

35 TGGTTTACCC AACAACATTG TGGATGTCTG GGAGTTTTTG GCCAAACCTA AACATTGCCA
900

GTATACATGG GATACACAAA TGAACCTCTAA TCTTGAATA ACTGCTGCTT GTAACTTCG
960

40 TTTTGATCGA ATATTTTCA GAGCAGCAGC AGAAGAGGGA CACATTATTC CCCGAAGTTT
1020

GGACCTTCTT GGATTAGAAA AACTGGACTG TGGTAGATTT CCTAGTGATC ACTGGGGTCT
1080

45 TCTGTGCAAC TTAGATATAA TATTGTAAAA TGCTTTTCAA GTGTGGGTTT TGCCCTGATT
1140

50 GTTGCAAATA CAATTTCCAC CTTCTGAAA GGTAGGTTTG CTGTGGAGGA AATAATGTAC
1200

TAGATCATTG TCACAGAAAA ACCAACTATG ATTTATGGTT GTGTTTTAG AATTCAACAT
1260

55 TAAAGATTAA TGTTTATTTA AACGAACACA TTCCTGCATT CAGGATGTGA GGCCATTAA
1320

TAAAAAGGGC ACAAAGCCTG TCAGAGTTTT CAACGGTGCT TACAGCTGCC AGCTGGATTG
 1380
 CAAACAGSTA CCCCATTTGTC TCTGAGCTAA TGTTTATATT TTTCATTCA GGCACCGAAA
 5 1440
 TAGTTAATAT TTAAATAAG TCTTCAAAG AAAACATAAG AGATTATTGA GTTCTTGGA
 1500
 CTGGATCCTT TATTTTATAA GTTCAGATCA TCTTAAATGA AAATGCCATG ATTATCTGCA
 10 1560
 GTTAAGTAGA TGACAGCTAT TCTACATCAG ACTTGATTTT TGTCAGCTAA TTACATAATT
 1620
 GGTAAGNTAT AATTGAAACC TTATGGCTTA AAATTCCTTA ACTCCTTTTT GATTCATGTT
 1680
 TGTAGTCATG TTGTCAACAG AGGCAAAGTT AAGCTTGATG ATGGTTAAAA TCGGTTTGAT
 20 1740
 AGCACCATGG GACATTTTTT TAACAAAAT AAATGCATGA AGAGACATAG CCTTTTAGTT
 1800
 TTGCTAATTG TGAAATGGAA ATGCTTTACA GGAAGTAAAT GCAAATTANT TTTAAGTGTG
 25 1860
 CTTTAAAGAA AAATATTTTC CCCACAGGAG AAATTTAAAT AAAGAATTTT ATTTGGTAAA
 1920

30

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:
 35 (A) LENGTH: 362 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

45 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Glu Leu Gly Ser Cys Leu Glu Gly Gly Arg Glu Ala Ala Glu Glu
 1 5 10 15
 Glu Gly Glu Pro Glu Val Lys Lys Arg Arg Leu Leu Cys Val Glu Phe
 20 25 30

Printed:22-04-2002

+32 9 2 SPEC

98201392

Ala Ser Val Ala Ser Cys Asp Ala Ala Val Ala Gln Cys Phe Leu Ala
35 40 45

5 Glu Asn Asp Trp Glu M t Glu Arg Ala Leu Asn Ser Tyr Phe Glu Pro
50 55 60

Pro Val Glu Glu Ser Ala Leu Glu Arg Arg Pro Glu Thr Ile Ser Glu
65 70 75 80

10 Pro Lys Thr Tyr Val Asp Leu Thr Asn Glu Glu Thr Thr Asp Ser Thr
85 90 95

15 Thr Ser Lys Ile Ser Pro Ser Glu Asp Thr Gln Gln Glu Asn Gly Ser
100 105 110

Met Phe Ser Leu Ile Thr Trp Asn Ile Asp Gly Leu Asp Leu Asn Asn
115 120 125

20 Leu Ser Glu Arg Ala Arg Gly Val Cys Ser Tyr Leu Ala Leu Tyr Ser
130 135 140

Pro Asp Val Ile Phe Leu Gln Glu Val Ile Pro Pro Tyr Tyr Ser Tyr
145 150 155 160

25 Leu Lys Lys Arg Ser Ser Asn Tyr Glu Ile Ile Thr Gly His Glu Glu
165 170 175

Gly Tyr Phe Thr Ala Ile Met Leu Lys Lys Ser Arg Val Lys Leu Lys
180 185 190

30 Ser Gln Glu Ile Ile Pro Phe Pro Ser Thr Lys Met Met Arg Asn Leu
195 200 205

35 Leu Cys Val His Val Asn Val Ser Gly Asn Glu Leu Cys Leu Met Thr
210 215 220

Ser His Leu Glu Ser Thr Arg Gly His Ala Ala Glu Arg Met Asn Gln
225 230 235 240

40 Leu Lys Met Val Leu Lys Lys Met Gln Glu Ala Pro Glu Ser Ala Thr
245 250 255

45 Val Ile Phe Ala Gly Asp Thr Asn Leu Arg Asp Arg Glu Val Thr Arg
260 265 270

Cys Gly Gly Leu Pro Asn Asn Ile Val Asp Val Trp Glu Phe Leu Gly
275 280 285

50 Lys Pro Lys His Cys Gln Tyr Thr Trp Asp Thr Gln Met Asn Ser Asn
290 295 300

Leu Gly Ile Thr Ala Ala Cys Lys Leu Arg Phe Asp Arg Ile Phe Phe
305 310 315 320

55 Arg Ala Ala Ala Glu Glu Gly His Ile Ile Pro Arg Ser Leu Asp Leu
325 330 335

Leu Gly Leu Glu Lys L u Asp Cys Gly Arg Phe Pro Ser Asp His Trp

Printed:22-04-2002

+32 9 2 SPEC

98201392

340

345

350

Gly Leu Leu Cys Asn Leu Asp Ile Ile Leu
355 360

5

(2) INFORMATION FOR SEQ ID NO: 3;

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 1312 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mus musculus

25

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:122..1234

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

30 AGCTATTAAT GATTGGAATT TATACGACTC ACTATAGGGA ATTTGGCCCT CGAGGCCAAG
60

AATTCGGCAC GAGGGCGGGA AGCAGCGTGA AGAGCGGCTG TTTTGAGGGG ACCCTGCGGC
120

35

G ATG GCG TCT GGC AGC AGT TCC GAT GCG GCG GAG CCC GCA GGG CCG
166

Met Ala Ser Gly Ser Ser Ser Asp Ala Ala Glu Pro Ala Gly Pro
1 5 10 15

40

GCA GGG CGG GCG GCG TCG GCG CCC GAA GCA GCA CAG GCG GAG GAG GAC
214

Ala Gly Arg Ala Ala Ser Ala Pro Glu Ala Ala Gln Ala Glu Glu Asp
20 25 30

45

CGG GTG AAG AGG CGG CGG CTT CAG TGC CTG GGC TTT GCG TTG GTG GGG
262

Arg Val Lys Arg Arg Arg Leu Gln Cys Leu Gly Phe Ala Leu Val Gly
35 40 45

50

GGA TGC GAC CCC ACG ATG GTC CCC AGC GTC CTG CGG GAG AAC GAC TGG
310

Gly Cys Asp Pro Thr Met Val Pro Ser Val Leu Arg Glu Asn Asp Trp
50 55 60

55

CAG ACG CAG AAA GCC CTG AGC GCC TAC TTC GAG CTG CCA GAG AAC GAC
358

Gln Thr Gln Lys Ala Leu Ser Ala Tyr Phe Glu Leu Pro Glu Asn Asp

Printed:22-04-2002

+32 9 2 SPEC

98201392

	65	70	75
	CAA GGG TGG CCG CGC CAG CCT CCC ACG TCC TTC AAG TCC GAG GCC TAT		
	406		
5	Gln Gly Trp Pro Arg Gln Pro Pro Thr Ser Phe Lys Ser Glu Ala Tyr		
	80	85	90 95
	GTT GAT CTA ACC AAC GAG GAT GCA AAT GAT ACA ACC ATT TTA GAA GCC		
	454		
10	Val Asp Leu Thr Asn Glu Asp Ala Asn Asp Thr Thr Ile Leu Glu Ala		
		100	105 110
	AGT CCA TCT GGA ACT CCT CTA GAA GAT AGC AGC ACT ATT TCT TTC ATT		
	502		
15	Ser Pro Ser Gly Thr Pro Leu Glu Asp Ser Ser Thr Ile Ser Phe Ile		
		115	120 125
	ACC TGG AAT ATT GAT GGA TTA GAT GGA TGC AAT CTG CCC GAG AGG GCT		
	550		
20	Thr Trp Asn Ile Asp Gly Leu Asp Gly Cys Asn Leu Pro Glu Arg Ala		
		130	135 140
	CGA GGG GTG TGT TCC TGC CTA GCT TTG TAT AGT CCA GAT GTG GTA TTT		
	598		
25	Arg Gly Val Cys Ser Cys Leu Ala Leu Tyr Ser Pro Asp Val Val Phe		
		145	150 155
	CTA CAG GAA GTT ATC CCC CCA TAC TGT GCC TAC CTA AAG AAG AGA GCA		
	646		
30	Leu Gln Glu Val Ile Pro Pro Tyr Cys Ala Tyr Leu Lys Lys Arg Ala		
		160	165 170 175
	GCC AGT TAC ACA ATT ATT ACA GGT AAT GAA GAA GGA TAT TTC ACA GCT		
	694		
35	Ala Ser Tyr Thr Ile Ile Thr Gly Asn Glu Glu Gly Tyr Phe Thr Ala		
		180	185 190
	ATA CTA TTG AAG AAA GGA AGA GTG AAA TTT AAA AGT CAG GAG ATT ATT		
	742		
40	Ile Leu Leu Lys Lys Gly Arg Val Lys Phe Lys Ser Gln Glu Ile Ile		
		195	200 205
	CCT TTT CCA AAT ACC AAA ATG ATG AGA AAC CTG CTA TGC GTA AAT GTG		
	790		
45	Pro Phe Pro Asn Thr Lys Met Met Arg Asn Leu Leu Cys Val Asn Val		
		210	215 220
	AGT TTG GGT GGA AAT GAA TTT TGC CTT ATG ACA TCC CAT TTG GAG AGC		
	838		
50	Ser Leu Gly Gly Asn Glu Phe Cys Leu Met Thr Ser His Leu Glu Ser		
		225	230 235
	ACC AGA GAA CAT TCT GCG GAA CGA ATA AGA CAA TTA AAA ACT GTT CTT		
	886		
55	Thr Arg Glu His Ser Ala Glu Arg Il Arg Gln Leu Lys Thr Val Leu		
		240	245 250 255

GGA AAA ATG CAA GAG GCT CCA GAT TCA ACC ACG GTT ATA TTT GCA GGA
 934
 Gly Lys Met Gln Glu Ala Pro Asp Ser Thr Thr Val Ile Phe Ala Gly
 260 265 270
 5
 GAT ACA AAT TTA AGA GAT CAA GAA GTT ATC AAA TGT GGT GGT TTA CCT
 982
 Asp Thr Asn Leu Arg Asp Gln Glu Val Ile Lys Cys Gly Gly Leu Pro
 275 280 285
 10
 GAC AAC GTT TTT GAT GCC TGG GAA TTT TTA GGC AAA CCT AAA CAT TGC
 1030
 Asp Asn Val Phe Asp Ala Trp Glu Phe Leu Gly Lys Pro Lys His Cys
 290 295 300
 15
 CAG TAT ACA TGG GAT ACG AAA GCA AAT AAC AAC CTC AGG ATC CCT GCT
 1078
 Gln Tyr Thr Trp Asp Thr Lys Ala Asn Asn Asn Leu Arg Ile Pro Ala
 305 310 315
 20
 GCT TAT AAG CAT CGT TTT GAT CGA ATA TTT TTC AGA GCA GAA GAG GGG
 1126
 Ala Tyr Lys His Arg Phe Asp Arg Ile Phe Phe Arg Ala Glu Glu Gly
 320 325 330 335
 25
 CAC CTT ATT CCT CAA AGT TTA GAC CTT GTT GGG TTG GAA AAA CTG GAC
 1174
 His Leu Ile Pro Gln Ser Leu Asp Leu Val Gly Leu Glu Lys Leu Asp
 340 345 350
 30
 TGT GGT AGA TTT CCG AGT GAT CAC TGG GGG CTC TTG TGC ACC TTG AAT
 1222
 Cys Gly Arg Phe Pro Ser Asp His Trp Gly Leu Leu Cys Thr Leu Asn
 355 360 365
 35
 GTA GTA TTG TGA AAAGCTTCCC ACTTGCAGCT TTACACGTTT GTTAGCACTA
 1274
 Val Val Leu *
 370
 40
 GTTCTGAATT TGTGTAGGTC TCAACCTTTC AGGACATC
 1312

45 (2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 371 amino acids

(B) TYPE: amino acid

50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

55 M t Ala Ser Gly Ser Ser Ser Asp Ala Ala Glu Pro Ala Gly Pro Ala
 1 5 10 15

Gly Arg Ala Ala Ser Ala Pro Glu Ala Ala Gln Ala Glu Glu Asp Arg

	20	25	30
	Val Lys Arg Arg Arg Leu Gln Cys Leu Gly Phe Ala Leu Val Gly Gly		
	35	40	45
5	Cys Asp Pro Thr Met Val Pro Ser Val Leu Arg Glu Asn Asp Trp Gln		
	50	55	60
10	Thr Gln Lys Ala Leu Ser Ala Tyr Phe Glu Leu Pro Glu Asn Asp Gln		
	65	70	75 80
	Gly Trp Pro Arg Gln Pro Pro Thr Ser Phe Lys Ser Glu Ala Tyr Val		
	85	90	95
15	Asp Leu Thr Asn Glu Asp Ala Asn Asp Thr Thr Ile Leu Glu Ala Ser		
	100	105	110
	Pro Ser Gly Thr Pro Leu Glu Asp Ser Ser Thr Ile Ser Phe Ile Thr		
	115	120	125
20	Trp Asn Ile Asp Gly Leu Asp Gly Cys Asn Leu Pro Glu Arg Ala Arg		
	130	135	140
	Gly Val Cys Ser Cys Leu Ala Leu Tyr Ser Pro Asp Val Val Phe Leu		
25	145	150	155 160
	Gln Glu Val Ile Pro Pro Tyr Cys Ala Tyr Leu Lys Lys Arg Ala Ala		
	165	170	175
30	Ser Tyr Thr Ile Ile Thr Gly Asn Glu Glu Gly Tyr Phe Thr Ala Ile		
	180	185	190
	Leu Leu Lys Lys Gly Arg Val Lys Phe Lys Ser Gln Glu Ile Ile Pro		
	195	200	205
35	Phe Pro Asn Thr Lys Met Met Arg Asn Leu Leu Cys Val Asn Val Ser		
	210	215	220
	Leu Gly Gly Asn Glu Phe Cys Leu Met Thr Ser His Leu Glu Ser Thr		
40	225	230	235 240
	Arg Glu His Ser Ala Glu Arg Ile Arg Gln Leu Lys Thr Val Leu Gly		
	245	250	255
45	Lys Met Gln Glu Ala Pro Asp Ser Thr Thr Val Ile Phe Ala Gly Asp		
	260	265	270
	Thr Asn Leu Arg Asp Gln Glu Val Ile Lys Cys Gly Gly Leu Pro Asp		
	275	280	285
50	Asn Val Phe Asp Ala Trp Glu Phe Leu Gly Lys Pro Lys His Cys Gln		
	290	295	300
	Tyr Thr Trp Asp Thr Lys Ala Asn Asn Asn Leu Arg Ile Pro Ala Ala		
55	305	310	315 320
	Tyr Lys His Arg Phe Asp Arg Ile Phe Phe Arg Ala Glu Glu Gly His		
	325	330	335

L u Ile Pro Gln Ser L u Asp Leu Val Gly Leu Glu Lys Leu Asp Cys
340 345 350

5 Gly Arg Phe Pro Ser Asp His Trp Gly Leu Leu Cys Thr Leu Asn Val
355 360 365

Val Leu *
370

10

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 1536 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

30

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 209..1536

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

35 AGAGAAAGAG GCTCCGGGGA GATAGCGGAC CAGTGAGGGC TGCCCCCTCTT TTGAAGCGGT
60

TTTCGTCTCT TTCCGCCAGT GGCCTCCCA G CTCACGCAGG GGCGGGTCCC GGTAGCGCGA
120

40

GGCGGTGCAG GGCGGGAAGG GGAGTGGTGG CGGCTGCGGC AGTAGGGACA GCAGGAGCAG
180

45

TGGTGCTGTC AGCGCGGCCG TCGGAGACAT GGGAGACCCG GGGTCGGAAA TAATAGAATC
240

TGTCCCTCCA GCTGGCCCTG AGGCATCTGA GTCAACAACG GATGAAAATG AAGACGACAT
300

50

TCAGTTTGTC AGTGAAGGAC CATCGAGACC TGTCTTTGAA TACATCGATC TGGTCTGTGG
360

TGATGATGAA AACCTAGCG CCTATTATAG TGATATTCTG TTTCCTAAAA TGCCAAAACG
420

55

ACAGGGTGAT TTTTGCATT TTTTAAATAT GAAGAAGGTG AAAACAGACA CAGAAAATAA
480

TGAAGTGAGC AAAATCACT GCAGATTGTC TAAGGCAAAG GAACCACATT TCGAGTATAT
 540
 5 AGAACAAACCA ATCATTGAAG AAAAGCCATC ACTTTCATCA AAGAAAGAAA TAGATAATCT
 600
 TGTGCTTCCA GATTGTTGGA ATGAAAAACA AGCATTATG TTTACAGAAC AATACAAATG
 660
 10 GCTTGAAATA AAAGAAGGTA AATTAGGATG TAAGGATTGT TCAGCAGTTC GGCATTGTTGGG
 720
 ATCGAAAGCA GAAAAGCATG TCCATGTGTC CAAGGAATGG ATTGCATATT TAGTAACCCC
 780
 15 TAATGGCAGT AATAAACTA CTAGGCAAGC TTCTCTACGA AAAAAATTA GGAACATGA
 840
 TGTCTCTAAA GCCCATGGTA AAATTCAGGA TTTGTTAAG GAATCAACTA ATGATTCAAT
 900
 20 TTGTAATTTA GTGCATAAAC AAAATAATAA AAATATTGAT GCTACTGTAA AAGTTTTCAA
 960
 25 TACTGTTTAC AGTTTAGTAA AACATAACAG ACCTTTATCT GATATTGAGG GGGCAAGAGA
 1020
 ATTACAGGAA AAAAATGGAG AGGTAAATTG TTAAATACA CGTTACAGTG CAACAAGAAT
 1080
 30 AGCAGAACAT ATTGCAAAAG AAATGAAGAT GAAGATATTT AAGAATATTA TAGAAGAGAA
 1140
 TGCCAAAATC TGTATCATAA TTGATGAGGC ATCTACAGTT TCAAAGAAAA CCACCCTAGT
 1200
 35 GATTTATCTC CAGTGCACAA TTCAGTCAGC TCCTGCACCT GTTATGTTAT TTGTGGCTTT
 1260
 40 AAAAGAATTG GTGTCAACTA TAGCAGAGTG TATTGTCAAT ACATTATTGA CTACTTTAAA
 1320
 TGATTGTGGT TTTACAAATG AATATTTGAA AGCAAATTTA ATTGCATTTT GTTCTGATGG
 1380
 45 TGCTAATACA ANCTGGGAA GAAAGTCTGG AGTAGCTACA AAATTGTTAG AAAATTTTCC
 1440
 TGAAATCATC ATTTGGAACT GTTTAAATCA TCGATTACAA TTGTCACTTG ATGATTCTAT
 1500
 50 ATCCGAAATA AAACAAATTA ATCATTTAAN NTATAA
 1536
 55 (2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 442 amino acids

Printed:22-04-2002

+32 9 2 SPEC

98201392

(B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: protein
 (iii) HYPOTHETICAL: NO
 (iv) ANTI-SENSE: NO
 10 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: homo sapiens

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met	Gly	Asp	Pro	Gly	Ser	Glu	Ile	Ile	Glu	Ser	Val	Pro	Pro	Ala	Gly	1	5	10	15
Pro	Glu	Ala	Ser	Glu	Ser	Thr	Thr	Asp	Glu	Asn	Glu	Asp	Asp	Ile	Gln	20	25	30	
Phe	Val	Ser	Glu	Gly	Pro	Ser	Arg	Pro	Val	Leu	Glu	Tyr	Ile	Asp	Leu	25	35	40	45
Val	Cys	Gly	Asp	Asp	Glu	Asn	Pro	Ser	Ala	Tyr	Tyr	Ser	Asp	Ile	Leu	50	55	60	
Phe	Pro	Lys	Met	Pro	Lys	Arg	Gln	Gly	Asp	Phe	Leu	His	Phe	Leu	Asn	30	65	70	75
Met	Lys	Lys	Val	Lys	Thr	Asp	Thr	Glu	Asn	Asn	Glu	Val	Ser	Lys	Asn	85	90	95	
His	Cys	Arg	Leu	Ser	Lys	Ala	Lys	Glu	Pro	His	Phe	Glu	Tyr	Ile	Glu	35	100	105	110
Gln	Pro	Ile	Ile	Glu	Glu	Lys	Pro	Ser	Leu	Ser	Ser	Lys	Lys	Glu	Ile	40	115	120	125
Asp	Asn	Leu	Val	Leu	Pro	Asp	Cys	Trp	Asn	Glu	Lys	Gln	Ala	Phe	Met	130	135	140	
Phe	Thr	Glu	Gln	Tyr	Lys	Trp	Leu	Glu	Ile	Lys	Glu	Gly	Lys	Leu	Gly	45	145	150	155
Cys	Lys	Asp	Cys	Ser	Ala	Val	Arg	His	Leu	Gly	Ser	Lys	Ala	Glu	Lys	165	170	175	
His	Val	His	Val	Ser	Lys	Glu	Trp	Ile	Ala	Tyr	Leu	Val	Thr	Pro	Asn	50	180	185	190
Gly	Ser	Asn	Lys	Thr	Thr	Arg	Gln	Ala	Ser	Leu	Arg	Lys	Lys	Ile	Arg	55	195	200	205
Glu	His	Asp	Val	S	r	Lys	Ala	His	Gly	Lys	Ile	Gln	Asp	Leu	Leu	210	215	220	

Printed: 22-04-2002

+32 9 2 SPEC

98201392

	Glu Ser Thr Asn Asp Ser Il Cys Asn Leu Val His Lys Gln Asn Asn	
	225	230 235 240
5	Lys Asn Ile Asp Ala Thr Val Lys Val Phe Asn Thr Val Tyr Ser Leu	
		245 250 255
	Val Lys His Asn Arg Pro Leu Ser Asp Ile Glu Gly Ala Arg Glu Leu	
		260 265 270
10	Gln Glu Lys Asn Gly Glu Val Asn Cys Leu Asn Thr Arg Tyr Ser Ala	
		275 280 285
	Thr Arg Ile Ala Glu His Ile Ala Lys Glu Met Lys Met Lys Ile Phe	
15		290 295 300
	Lys Asn Ile Ile Glu Glu Asn Ala Lys Ile Cys Ile Ile Ile Asp Glu	
		305 310 315 320
20	Ala Ser Thr Val Ser Lys Lys Thr Thr Leu Val Ile Tyr Leu Gln Cys	
		325 330 335
	Thr Ile Gln Ser Ala Pro Ala Pro Val Met Leu Phe Val Ala Leu Lys	
		340 345 350
25	Glu Leu Val Ser Thr Ile Ala Glu Cys Ile Val Asn Thr Leu Leu Thr	
		355 360 365
	Thr Leu Asn Asp Cys Gly Phe Thr Asn Glu Tyr Leu Lys Ala Asn Leu	
30		370 375 380
	Ile Ala Phe Cys Ser Asp Gly Ala Asn Thr Xaa Leu Gly Arg Lys Ser	
		385 390 395 400
35	Gly Val Ala Thr Lys Leu Leu Glu Asn Phe Pro Glu Ile Ile Ile Trp	
		405 410 415
	Asn Cys Leu Asn His Arg Leu Gln Leu Ser Leu Asp Asp Ser Ile Ser	
		420 425 430
40	Glu Ile Lys Gln Ile Asn His Leu Xaa Tyr	
		435 440

CLAIMS

1. A functional protein, capable of interacting with the cytoplasmic domain of CD40, wherein said protein has no homology to TRAF-proteins.
- 5 2. A functional protein according to claim 1, comprising an amino acid sequence with 70-100% homology to the amino acid sequence depicted in SEQ ID NO. 2.
3. A functional protein according to claim 1, comprising an amino acid sequence with 70-100% homology to the amino acid sequence depicted in
10 SEQ ID NO. 4.
4. A functional protein according to claim 1, comprising an amino acid sequence with 70-100% homology to the amino acid sequence depicted in SEQ ID NO. 6.
5. A functional protein according to claim 1, comprising an amino acid
15 sequence with 70-100% homology to the amino acids 274-362 of SEQ ID NO. 2.
6. A functional protein according to claim 1, comprising an amino acid sequence with 70-100% homology to the amino acids 2-245 of SEQ ID NO.6.
- 20 7. A nucleic acid encoding a protein according to any of the claims 1-6.
8. A nucleic acid according to claim 7, with about 70-100% homology to the DNA sequence depicted in SEQ ID NO. 1.
9. A nucleic acid according to claim 7, with about 70-100% homology to the DNA sequence depicted in SEQ ID NO.3.

10. A nucleic acid according to claim 7, with about 70-100% homology to the DNA sequence depicted in SEQ ID NO. 5.
11. The use of a functional protein, according to any of the claims 1-6 and/or a functional fragment thereof to treat CD40-related diseases and/or NF- κ B related diseases.
12. The use according to claim 11 in which the disease is atherosclerosis, arthritis, multiple sclerosis, systemic lupus erythematosus and/or graft rejection.
13. The use of a functional protein according to any of the claims 1-6 and/or a functional fragment thereof to sensitise tumor cells to anti-tumor treatments.
14. The use of a functional protein according to any of the claims 1-6 and/or a functional fragment thereof to screen for compounds that interfere with the interaction of said protein(s) with other compounds of the CD40 or NF- κ B related pathway.
15. A method for screening compounds comprising the use of a protein according to claim 14.
16. A compound isolated with the method according to claim 15.
17. A pharmaceutical composition comprising one or more functional proteins according to any of the claims 1-6 and/or functional fragments thereof and a pharmaceutical acceptable carrier material.
18. A pharmaceutical composition comprising one or more compounds according to claim 16 and a pharmaceutical acceptable carrier material.

19. Use of a protein according to any of the claims 1-6 and/or functional fragments thereof for the manufacture of a pharmaceutical composition to treat CD40 and/or NF- κ B related diseases.

ABSTRACT

The present invention relates to novel proteins that interact with the cytoplasmic domain of CD40, which are useful in the treatment of CD40 and/or NF- κ B related diseases. Surprisingly, these proteins do not show significant homology with the TRAF-protein family, and offer therefore the possibility to modulate the CD40 and/or NF- κ B pathway independently from the TRAF-CD40 interaction.

Figure 1




Figure 2

